XFORCE Disinfectant is intended for use in: hospitals and healthcare settings, ambulances, anesthesia, CAT labs, CCU, Central Supply, Clinics, Dental Offices, Dialysis Clinics, Doctor's office(s), Emergency Rooms, ER, Home health care, Hospices, ICU, Laboratory, Laundry rooms, Neonatal Intensive Care Units (NICU), Newborn nurseries, Nursing homes, Operating rooms, Pediatrics, Pediatric Intensive Care Units (PICU), Physical therapy, Physician's office(s), Opthalmic offices, Radiology, Recovery Room, Rehabilitation, Respiratory therapy, Surgery, Transport vehicles, X-Ray, Critical Care Areas, XFORCE Disinfectant will clean and disinfect hard non-porous surfaces of: bed railings, cabinets, carts, chairs, counters, tables, telephones, toilet seats, toilets, bathtubs, bathroom fixtures, shower stalls, sinks, tiles and vanity tops. Disinfects in just 2 minutes. *One step cleaning and disinfecting. Kills Rhinovirus, causative agent of the common cold, in 2 minutes! FORCE disinfectant kills microorganisms on hospital soft surfaces, beddings, couches, curtains, and upholstery. A"One Step disinfectant" claim is limited to association with non-food contact surface use.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This product is not for use on critical and semi-critical medical device surfaces. GENERAL CLEANING: Spray soiled area then wipe with a dry paper towel or lint-free cloth. DISINFECTING, DEODORIZING: Remove gross filth or heavy soil prior to application of XFORCE Disinfectant, Apply product with a hand pump trigger sprayer. Hold sprayer six to eight inches from the surface to be treated. Spray area until it is covered with the product. Allow surface to remain visibly wet for 2 minutes. No scrubbing necessary. Allow treated surfaces to air dry or wipe off with a clean cloth. When using on food contact surfaces: counter tops, stovetops, highchairs, kitchen appliances, thoroughly rinse all treated surfaces with potable water. This product must not be used to clean eating utensils, glassware and dishes. XFORCE Disinfectant is an effective disinfectant against: Pseudomonas aeruginosa, Salmonella enterica (Salmonella). Staphylococcus aureus, Methicillin Resistant Staphylococcus aureus (MRSA), Vancomycin Resistant Enterococcus faecium, Multi-Drug Resistant (MDR) Acinetobacter baumannii Ceftazidime and Gentamicin Resistant, Extended Spectrum Beta-Lactamase (ESBL) Escherichia coli and Klebsiella pneumoniae New Dehli Metallo-Beta Lactamase (NDM-1) Carbapenem Resistant.



EPA Reg. No. 1839-220-70053 EP

EPA EST. No. 6574-KY-001

of Bacteria in 5 Seconds!** Disinfects household, kitchen and bathroom bacteria† Hospital disinfectant ready-to-use RTU spray

** Staphylococcus aureus (Staph), Klebsiella pneumoniae †Pseudomonas aeruginosa, Staphylococcus aureus (Staph), Salmonella enterica (Salmonella), Methicillin Resistant Staphylococcus aureus (MRSA), Vancomycin Resistant Enterococcus faecium (VRE), Multi-Drug Resistant (MDR) Acinetobacter baumannii Ceftazidime and Gentamicin Resistant, Extended Spectrum Beta-Lactamase Escherichia coli (E.coli), Klebsiella pneumoniae New Delhi Metallo-Beta Lactamase (NDM-1) Carbapenem Resistant

ACTIVE INGREDIENTS:

ACTIVE INGREDIENTS:	
Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈)	
dimethyl benzyl ammonium chlorides	0.15%
Alkyl (68% C ₁₂ , 32% C ₁₄) dimethyl	
ethylbenzyl ammonium chlorides	0.15%
INERT INGREDIENTS:	99.70%
TOTAL	100 00%

KEEP OUT OF REACH OF CHILDREN

See side panels for additional precautionary statements.

NET CONTENTS: 1 QT. (0.946 LITER) VIRUCIDAL ACTIVITY: XFORCE Disinfectant is an effective virucide on hard non-porous surfaces against Avian Influenza Virus (H5N1) NIBRG-14, when the treated surface is allowed to remain wet for 30 seconds. XFORCE Disinfectant is an effective virucide on hard, non-porous surfaces against Human Rotavirus (ATCC VR-2018) and Bovine Viral Diarrhea Virus (BVDV) when the treated surface is allowed to remain wet for 1 minute. XFORCE Disinfectant is an effective virucide on hard, non-porous surfaces against Rhinovirus Type 14 and Rhinovirus Type 39 when treated surface is allowed to remain wet for 2 minutes. Allow treated surfaces to air dry or wine off with a clean cloth.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. IF IN EYES: Hold eyes open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice. IF ON SKIN: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.

PESTICIDE STORAGE - Store in a dry place inaccessible to children.

container. Do not reuse or refill this container. Offer for recycling if available or puncture and dispose in a sanitary landfill, incineration, or burning, if allowed by state and local authorities. If burned, stay out of smoke. PESTICIDE DISPOSAL - Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for quidance.

108699-L 1QT 5/27/20

